



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search PubMed		▼ for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details		

- The Clipboard will hold a maximum of 500 items.
- Clipboard items will be lost after one hour of inactivity.

Entrez
PubMed

Display	Summary	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
Items 1-15 of 15									One page.

PubMed
Services

☐ 1: [Kotenko SV, Izotova LS, Mirochnitchenko OV, Esterova E, Dickensheets H, Donnelly RP, Pestka S.](#) Related Articles, Links

Identification, cloning, and characterization of a novel soluble receptor that binds IL-22 and neutralizes its activity.

J Immunol. 2001 Jun 15;166(12):7096-103.

PMID: 11390454 [PubMed - indexed for MEDLINE]

☐ 2: [Kotenko SV, Izotova LS, Mirochnitchenko OV, Esterova E, Dickensheets H, Donnelly RP, Pestka S.](#) Related Articles, Links

Identification of the functional interleukin-22 (IL-22) receptor complex: the IL-10R2 chain (IL-10Rbeta) is a common chain of both the IL-10 and IL-22 (IL-10-related T cell-derived inducible factor, IL-TIF) receptor complexes.

J Biol Chem. 2001 Jan 26;276(4):2725-32.

PMID: 11035029 [PubMed - indexed for MEDLINE]

☐ 3: [Ding Y, Qin L, Kotenko SV, Pestka S, Bromberg JS.](#) Related Articles, Links

A single amino acid determines the immunostimulatory activity of interleukin 10.

J Exp Med. 2000 Jan 17;191(2):213-24.

PMID: 10637267 [PubMed - indexed for MEDLINE]

☐ 4: [Gruenberg BH, Schoenemeyer A, Weiss B, Toschi L, Kunz S, Wolk K, Asadullah K, Sabat R.](#) Related Articles, Links

A novel, soluble homologue of the human IL-10 receptor with preferential expression in placenta.

Genes Immun. 2001 Oct;2(6):329-34.

PMID: 11607789 [PubMed - indexed for MEDLINE]

☐ 5: [Xu W, Presnell SR, Parrish-Novak J, Kindsvogel W, Jaspers S, Chen Z, Dillon SR, Gao Z, Gilbert T, Madden K, Schlusmeyer S, Yao L, Whitmore TE, Chandrasekhar Y, Grant FJ, Maurer M, Jelinek L, Storey H, Brender T, Hammond A, Topouzis S, Clegg CH, Foster DC.](#) Related Articles, Links

A soluble class II cytokine receptor, IL-22RA2, is a naturally occurring IL-22 antagonist.

Proc Natl Acad Sci U S A. 2001 Aug 14;98(17):9511-6.

PMID: 11481447 [PubMed - indexed for MEDLINE]

Related
Resources

- ☐ 6: Xie MH, Aggarwal S, Ho WH, Foster J, Zhang Z, Stinson J, Wood WI, Goddard AD, Gurney AL. Related Articles, Links
Interleukin (IL)-22, a novel human cytokine that signals through the interferon receptor-related proteins CRF2-4 and IL-22R.
J Biol Chem. 2000 Oct 6;275(40):31335-9.
PMID: 10875937 [PubMed - indexed for MEDLINE]
- ☐ 7: Dumoutier L, Lejeune D, Colau D, Renault JC. Related Articles, Links
Cloning and characterization of IL-22 binding protein, a natural antagonist of IL-10-related T cell-derived inducible factor/IL-22.
J Immunol. 2001 Jun 15;166(12):7090-5.
PMID: 11390453 [PubMed - indexed for MEDLINE]
- ☐ 8: Dumoutier L, Van Roost E, Ameye G, Michaux L, Renault JC. Related Articles, Links
IL-TIF/IL-22: genomic organization and mapping of the human and mouse genes.
Genes Immun. 2000 Dec;1(8):488-94.
PMID: 11197690 [PubMed - indexed for MEDLINE]
- ☐ 9: Dumoutier L, Van Roost E, Colau D, Renault JC. Related Articles, Links
Human interleukin-10-related T cell-derived inducible factor: molecular cloning and functional characterization as an hepatocyte-stimulating factor.
Proc Natl Acad Sci U S A. 2000 Aug 29;97(18):10144-9.
PMID: 10954742 [PubMed - indexed for MEDLINE]
- ☐ 10: Dumoutier L, Louahed J, Renault JC. Related Articles, Links
Cloning and characterization of IL-10-related T cell-derived inducible factor (IL-TIF), a novel cytokine structurally related to IL-10 and inducible by IL-9.
J Immunol. 2000 Feb 15;164(4):1814-9.
PMID: 10657629 [PubMed - indexed for MEDLINE]
- ☐ 11: Tachiiri A, Imamura R, Wang Y, Fukui M, Umemura M, Suda T. Related Articles, Links
Genomic structure and inducible expression of the IL-22 receptor alpha chain in mice.
Genes Immun. 2003 Mar;4(2):153-9.
PMID: 12618864 [PubMed - in process]
- ☐ 12: Logsdon NJ, Jones BC, Josephson K, Cook J, Walter MR. Related Articles, Links
Comparison of interleukin-22 and interleukin-10 soluble receptor complexes.
J Interferon Cytokine Res. 2002 Nov;22(11):1099-112.
PMID: 12513909 [PubMed - in process]
- ☐ 13: Kotenko SV. Related Articles, Links
The family of IL-10-related cytokines and their receptors: related, but to what extent?
Cytokine Growth Factor Rev. 2002 Jun;13(3):223-40. Review.
PMID: 12486876 [PubMed - indexed for MEDLINE]
- ☐ 14: Nagem RA, Colau D, Dumoutier L, Renault JC, Ogata C, Polikarpov I. Related Articles, Links
Crystal structure of recombinant human interleukin-22.
Structure (Camb). 2002 Aug;10(8):1051-62.
PMID: 12176383 [PubMed - indexed for MEDLINE]

- ☐ 15: [Spencer SD, Di Marco F, Hooley J, Pitts-Meek S, Bauer M, Ryan AM, Sordat B, Gibbs VC, Aguet M.](#) Related Articles, Links



The orphan receptor CRF2-4 is an essential subunit of the interleukin 10 receptor.
J Exp Med. 1998 Feb 16;187(4):571-8.
PMID: 9463407 [PubMed - indexed for MEDLINE]

Display	Summary	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
Items 1-15 of 15									One page.

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)